arXiv:astro-ph/9803025v1 3 Mar 1998

RXTE/ASM OBSERVATIONS OF PSR B1259−63

Victoria M. KASPI and Ronald A. REMILLARD
Department of Physics and Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
vicky@space.mit.edu, rr@space.mit.edu

Abstract

RXTE/ASM observations of PSR B1259−63 near the 1997 periastron passage find no emission brighter than ∼10 times that detected by *ASCA* during the 1994 periastron encounter, demonstrating that the pulsar is unlikely to have undergone even brief episodes of accretion.

1. Introduction, Observations, and Conclusions

The 48 ms radio pulsar PSR B1259−63 is in a highly eccentric 3.4 yr orbit with a Be star (Johnston et al. 1992). Near periastron, for a sufficiently strong Be star wind, PSR B1259−63 could possibly make a transition to an accretion regime, exhibiting X-ray pulsations of luminosity $>10^{35}$ erg s^{-1}. *ASCA* X-ray observations at three epochs around the 1994 periastron passage (Kaspi et al. 1994; Hirayama 1996) found the source to be unpulsed, with modest luminosity $\sim 10^{34}$ erg s^{-1}, for a distance of 2 kpc. These properties imply that shock emission produces the observed X-rays, rather than accretion (Tavani & Arons 1997). However radio timing observations suggested sudden, brief accretion events near periastron that could not be ruled out by the X-ray observations (Manchester et al. 1995). Indeed the pulsar's anomalously low period suggests it may have undergone occasional accretion episodes in the past. *ASCA* observations during the 1997 periastron passage were impossible due to solar constraints.

The ASM on *RXTE* has been observing bright celestial X-ray sources regularly since early 1996. The instrument consists of three "scanning shadow cameras," each containing a position-sensitive proportional counter that is mounted below a wide-field collimator, covered by a coded mask. The instrument provides roughly 5 celestial scans per day, with diminished exposure in directions toward the Sun. Further information on the ASM is given by Levine et al. (1996). While



Fig. 1. (a) ASM light curve for PSR B1259−63. (b) 3σ upper limits to the ASM flux near periastron from (a). The horizontal dashed line represents the brightest *ASCA* 1994 periastron detection (Kaspi et al. 1994), and the vertical dotted line indicates periastron.

PSR B1259−63 is not routinely monitored for the ASM source-history database, we extracted an X-ray light curve as an archival analysis project using standard techniques. The derived light curve (Figure 1a) covers the time interval of MJD 50178–50762 (1996 Apr 5 to 1997 Nov 10). The overall mean intensity of PSR B1259−63 during this time interval was $(0.5 \pm 2.4) \times 10^{-11}$ erg s^{-1} cm^{-2} at 2–10 keV. This result is given with consideration of both the systematic bias and uncertainty in the ASM light curves for faint, yet fairly isolated X-ray sources. Figure 1b shows 3σ upper limits obtained in the periastron vicinity in 2-day averages. It shows that in the 2 weeks following periastron, the source was not observed to be brighter than ∼10 times the 1994 periastron luminosity.

The 3σ upper limits to the X-ray emission from PSR B1259−63 from the ASM thus argue that the pulsar did not undergo even brief episodes of accretion during the 1997 periastron passage. This is consistent with the shock emission model, as well as with new conclusions based on timing that rule out the sudden "spin-ups" that were claimed previously (Wex et al. 1998).

2. References

Hirayama, M. 1996, PhD Thesis, University of Tokyo
Kaspi, V. M. et al. 1995, ApJ, 453, 424
Levine, A. M. et al. 1996, ApJ, 469, L33
Manchester, R. N. et al. 1995, ApJ, 445, L137
Tavani, M., Arons, J. 1997, ApJ, 477, 439
Wex, N. et al. 1998, MNRAS, in press